



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification: G06F 19/00	A2	(11) International Publication Number: WO 00/43942 (43) International Publication Date: 27 July 2000 (27.07.2000)
(21) International Application Number: PCT/US00/02000 (22) International Filing Date: 25 January 2000 (25.01.2000) (30) Priority Data: 60/116,956 25 January 1999 (25.01.1999) US (60) Parent Application or Grant COMBIMATRIX CORPORATION [/]; (). ANDERSON, Brooke, P. [/]; (). MONTGOMERY, Donald, D. [/]; (). HALLUIN, Albert, P. ; ().	Published 09/490, 563	
(54) Title: A METHOD FOR SELECTING OLIGONUCLEOTIDES HAVING LOW CROSS HYBRIDIZATION (54) Titre: TECHNIQUE PERMETTANT DE SELECTIONNER DES OLIGONUCLEOTIDES A FAIBLE HYBRIDATION CROISEE		
(57) Abstract <p>The present invention provides methods to provide a set of probes that hybridize or bind relatively well to their intended targets but that do not substantially bind or hybridize to unintended targets. The quality of hybridizing or binding relatively well to intended targets but not to unintended targets may be quantified using delta Tm according to the methods of the present invention. The present invention also provides sets of probes produced according to the subject methods and programmed computer systems functional to perform the subjects methods.</p> (57) Abrégé <p>L'invention concerne des techniques permettant de fournir un ensemble de sondes qui s'hybrident ou se lient relativement bien aux cibles souhaitées, mais qui s'hybrident ou se lient sensiblement aux cibles non souhaitées. La qualité de l'hybridation ou de la liaison peut être quantifiée à l'aide de delta Tm en fonction des techniques de l'invention. Cette invention fournit également des ensembles de sondes produites en fonction desdites techniques, et des systèmes informatiques programmés fonctionnellement de façon à utiliser les techniques décrites.</p>		

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : G06F 19/00	A2	(11) International Publication Number: WO 00/43942 (43) International Publication Date: 27 July 2000 (27.07.00)
(21) International Application Number: PCT/US00/02000 (22) International Filing Date: 25 January 2000 (25.01.00) (30) Priority Data: 60/116,956 25 January 1999 (25.01.99) US (71) Applicant: COMBIMATRIX CORPORATION [US/US]; Suite 210, 887 Mitten Road, Burlingame, CA 94010 (US). (72) Inventors: ANDERSON, Brooke, P.; 1410 Millbrae Avenue #105, Millbrae, CA 94030 (US). MONTGOMERY, Donald, D.; 1410 Millbrae Avenue #306, Millbrae, CA 94030 (US). (74) Agent: HALLUIN, Albert, P.; Howrey & Simon, 1299 Pennsylvania Avenue, N.W., Washington, DC 20004 (US).		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>Without international search report and to be republished upon receipt of that report.</i>
(54) Title: A METHOD FOR SELECTING OLIGONUCLEOTIDES HAVING LOW CROSS HYBRIDIZATION		
(57) Abstract		
<p>The present invention provides methods to provide a set of probes that hybridize or bind relatively well to their intended targets but that do not substantially bind or hybridize to unintended targets. The quality of hybridizing or binding relatively well to intended targets but not to unintended targets may be quantified using delta T_m according to the methods of the present invention. The present invention also provides sets of probes produced according to the subject methods and programmed computer systems functional to perform the subjects methods.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LJ	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

Description

5

10

15

20

25

30

35

40

45

50

55

**A METHOD FOR SELECTING OLIGONUCLEOTIDES
HAVING LOW CROSS HYBRIDIZATION**

FIELD OF THE INVENTION

The present invention is in the field of biological and chemical synthesis and processing. The present invention relates to methods for selecting oligonucleotides for low cross hybridization. The present invention may be applied in the field of, but is not limited to, the field of chemical or biological synthesis, diagnostics and therapeutics.

The present application claims priority to U.S. Provisional Patent Application Serial No. 60/116,956 filed January 25, 1999.

BACKGROUND OF THE INVENTION

Advances are emerging continually in the field of biological and chemical processing and synthesis. Many novel and improved solid phase arrays or "gene chips" are being developed providing rapid methods for synthesizing chemical and biological materials. Examples of such technologies include those described by Pirrung *et al.*, U.S. Patent No. 5,143,854, those described by Southern in WO 93/22480, those described by Heller in WO 95/12808 and those described by Montgomery in PCT/US97/11463. Hence, it is possible to synthesize, to manipulate and to examine ever increasing amounts of genetic materials. Moreover, it is possible to work simultaneously with, to analyze, and to test ever larger amounts of genetic materials.

Oligonucleotides can hybridize or bind to other oligonucleotides depending upon whether or not their sequences are more or less complementary. Sometimes, it is desirable to find a set of oligonucleotides that, as much as possible within a given set of constraints, do not hybridize or bind to each other. Optimization methods can be used to assist in selecting such a set of oligonucleotides.

There are many possibilities of how to make an oligonucleotide or oligonucleotides from a longer oligonucleotide. For example, one may cut the long oligonucleotide or select a segment or segments from which to form a smaller oligonucleotide. These smaller oligonucleotides formed by these and other methods known to those skilled in the art may be referred to as "substring sequences". There is great flexibility on which substring of the oligonucleotide to select as a target sequence for binding.

Mitsubishi *et al.*, U.S. Patent No. 5,556,749 describe a computerized method for designing optimal DNA probes. The method is intended to produce probes designed for diagnosis and monitoring. However, the method described therein does not contemplate choosing more than one probe for simultaneous use with another probe whereby interaction and cross hybridization is minimized.

5 It is an object of the present invention to provide a method for choosing complementary substrings from among target oligonucleotides such that the substrings bind relatively well to their target oligonucleotides but do not substantially bind to other oligonucleotides in a sample. For instance, given a long oligonucleotide (e.g., a string of RNA, mRNA, DNA or cDNA) it is possible to select a piece or pieces from it for a later experiment, e.g. as a capture probe. In the case of an immobilized capture probe, it is desirable that the oligonucleotide substring sequence bind to the long oligonucleotide that it came from but not to other oligonucleotides that might be present in a sample. It is often desirable to prepare an array of such immobilized capture probes so that each one binds or hybridizes to its intended target but does not bind or hybridize strongly to any of the other targets in the solution. An array of immobilized capture probes constructed according to the methods described allows using a smaller array of capture-probes for sequestering a desired set of targets because the array does not require as much redundancy for elucidating data points as other arrays where binding is not as clearly discernible.

SUMMARY OF THE INVENTION

25 In a first aspect, the present invention features methods to provide a set of probes that hybridize or bind relatively well to their intended targets but do not bind substantially to unintended targets. The quality of hybridizing or binding relatively well to intended targets but not to unintended targets may be quantified using delta Tm (difference in strength of hybridization, such as difference in melting temperatures) according to the methods of the present invention. A probe having a relatively small delta Tm generally hybridizes to at least one unintended target substantially well. A probe having a relatively large delta Tm has substantially no unintended targets that it hybridizes to substantially well.

35 The methods according to the present invention feature:

1. Determining a set of targets. In some circumstances where it is not clear what the identity of all the targets in a particular solution might be, it is possible to determine a list of some of the targets that might be in the particular solution and to include that list in the set of targets. If there are some targets in the list that are not actually in the solution, it does not harm the quality of probes selected according to the present methods.
2. Selecting a particular target from the set of targets. This becomes the current target.
3. Choosing a sequence substring from the current target and providing its complementary sequence. This is a candidate probe. Choosing a sequence substring may be done by starting at a particular point in the current target and then incrementing the starting point by some amount each time a new substring is chosen, with wraparound of the starting point to the front portion

5 of the current target when an increment would otherwise run off the end of the target. A substring
may be chosen at random or any arbitrary function may be applied in order to determine which
substring to choose. If there are no more substrings in the current target that have not already been
10 tried, it is recommended to: (i) use the best candidate probe selected so far noting that this target
might not be as selectively captured as desired, (ii) to do no more picking of probes for this target, and
(iii) to return to step 2, *supra*. Otherwise, using the candidate probe as determined and proceeding to
step 4, *infra*.

15 4. Determining whether the candidate probe satisfies any criteria required or desired for
the set of probes. If it does not meet such criteria, returning to step 3 and choose a new candidate
probe. If it does satisfy such criteria, proceeding to step 5.

20 5. Calculating the T_m for the candidate probe using the hybridization model.
Calculating substantially all possible cross T_m's of the candidate probe hybridizing to all unintended
targets and finding the maximum cross T_m. Calculating delta T_m. Note that the set of all unintended
targets will include previously picked probes if probes are also to be in solution as opposed to affixed
to a support.

25 6. Proceeding to step 7 if delta T_m is acceptably large. Returning to step 3 and choosing
a new candidate probe if delta T_m is not acceptable large.

30 7. At this point, a suitable probe has been chosen. The foregoing procedure may be
repeated additional times to choose other probes for additional targets or, if desired, additional probes
for a target for which one has already found one or more probes.

In a second aspect, the present invention features a set of probes that hybridize or bind
relatively well to their intended targets but do not bind substantially to unintended targets.

35 In a third aspect, the present invention features a set of probes that hybridize or bind relatively
well to their intended targets but do not bind substantially to unintended targets produced in
accordance with the methods set forth, *supra*.

40 In a fourth aspect, the present invention features a programmed computer system for providing
the sequences of a set of probes that hybridize or bind relatively well to their intended targets but do
not bind substantially to unintended targets.

45 DETAILED DESCRIPTION OF THE INVENTION

The present invention features methods to provide a set of probes that hybridize or bind
relatively well to their intended targets but do not bind substantially to unintended targets. The quality
of hybridizing or binding relatively well to intended targets but not to unintended targets may be

5 quantified using delta T_m according to the methods of the present invention. A probe with a small delta T_m generally hybridizes to at least one unintended target relatively well. A probe having a large delta T_m has substantially no unintended targets that it hybridizes to relatively well.

10 The methods according to the present invention feature:

1. Determining a set of targets. In some circumstances where it is not clear what the identity of all the targets in a particular solution might be, it is possible to determine a list of some of the targets that might be in the particular solution and to include that list in the set of targets. If there are some targets in the list that are not actually in the solution, it does not harm the quality of probes selected according to the present methods.

2. Selecting a particular target from the set of targets. This becomes the current target for which one desires a probe.

3. Choosing a sequence substring from the current target and providing its complementary sequence. This is a candidate probe. Choosing a sequence substring may be done by starting at a particular point in the current target and then incrementing the starting point by some amount each time a new substring is chosen, with wraparound of the starting point to the front portion of the current target when an increment would otherwise run off the end of the target. A substring may be chosen at random or any arbitrary function may be applied in order to determine which substring to choose. If there are no more substrings in the current target that have not already been tried, it is recommended to: (i) use the best candidate probe selected so far noting that this target might not be as selectively captured as desired, (ii) to do no more picking of probes for this target, and (iii) to return to step 2, *supra*. Otherwise, using the candidate probe as determined and proceeding to step 4, *infra*.

4. Determining whether the candidate probe satisfies any criteria required or desired for the set of probes. If it does not meet such criteria, returning to step 3 and choose a new candidate probe. If it does satisfy such criteria, proceeding to step 5.

5. Calculating the T_m for the candidate probe using the hybridization model. Calculating substantially all possible cross T_m's of the candidate probe hybridizing to all unintended targets and finding the maximum cross T_m. Calculating delta T_m. Note that the set of all unintended targets will include previously picked probes if probes are also to be in solution as opposed to affixed to a support.

6. Proceeding to step 7 if delta T_m is acceptably large. Returning to step 3 and choosing a new candidate probe if delta T_m is not acceptable large.

5

7. At this point, a suitable probe has been chosen. The foregoing procedure may be repeated additional times to choose other probes for additional targets or, if desired, additional probes for a target for which one has already found one or more probes.

10

In particular embodiments where the probes and targets are both in solution (*i.e.*, the probes are not tethered to a support), it is preferable to include each previously accepted probe into the list of targets before calculating delta T_m for the current candidate probe. Those skilled in the art will readily understand that it is preferable to do so because the probes are able to hybridize to each other as well as to the targets and so should be included in any calculation of the strength of unintended hybridizations. Those skilled in the art will readily understand that this feature means that the present invention provides probes having reduced cross hybridization with each other. Therefore, the present invention is particularly applicable to instances where multiple probes are used simultaneously in solution. This is often the case where primers are being used such as to amplify or copy sections of genetic material. In that case, "probe" and "primer" are meant to refer to the same oligonucleotide.

15

20

25

30

35

40

45

50

Adding targets as the methods according to the present invention progress may interfere with the ability to find a probe for a particular target not because of conflict with other targets but because of conflict (too strong a hybridization) to previously picked probes. If performing the methods outlined in the present invention reveals that there are a relatively large number of unacceptable probes found, and if these probes are unsuitable because they bind too strongly to other probes, it is generally preferred to begin the method from the start using a different order of picking current targets. This generally results in a different order of picking probes and can result in a set of probes that are more mutually exclusive. For instance, if the first time, probes for targets are picked in the following order: target 1, target 5, target 2, target 4, target 3, and the best probes for target 2 and target 3 do not have an acceptable delta T_m because they bind relatively well to probes for target 1 and target 5, it is possible to repeat the methods according to the present invention again by picking probes for targets in for example, the following order: target 2, target 3, target 5, target 4, target 1. In effect, the sequence for choosing targets may be modified within the present methods. Alternatively, it is possible to start with the set of probes found so far, look at the probe that was found to be less than ideal, find what other probes it hybridizes too strongly with, redo those probes, and repeat this process until a compatible set of probes is found. By way of example, suppose that a set of probes is found, but probes 2 and 5 do not have an acceptable delta T_m because they hybridize too strongly with probes 1 and 3, respectively. It is possible to eliminate probe 1 and proceed according to the methods of the present invention again for target 1 leaving the rest of the probes as is, finding a new acceptable probe for target 1 that perhaps does not conflict with probe 2. Then a skilled artisan may do the same for probe 3. This process (of redoing the previously found probes that later are found to conflict with other probes) may be iterated

55

5

until a good solution is found. If no suitable probes are found, one or more targets may be eliminated from the solution of interest.

10

In a second aspect, the present invention features a set of probes that hybridize or bind relatively well to their intended targets but do not bind substantially to unintended targets. The quality of hybridizing or binding relatively well to intended targets but not to unintended targets may be quantified using delta T_m . A probe having a small delta T_m generally hybridizes to at least one unintended target relatively well. A probe having a large delta T_m has substantially no unintended targets that it hybridizes to relatively well. Therefore, the set of probes according to the present invention may have, for example, a delta T_m of 5° C, 10° C or, for greater separation, 20° C. In general, the larger the delta T_m , the easier to dehybridize unintended targets while maintaining the intended targets hybridized by the probes in the subject set of probes.

20

In a third aspect, the present invention features a set of probes that hybridize or bind relatively well to their intended targets but do not bind substantially to unintended targets produced in accordance with the methods set forth, *supra*.

25

In a fourth aspect, the present invention features a programmed computer system for providing the sequences of a set of probes that hybridize or bind relatively well to their intended targets but do not bind substantially to unintended targets. Such a programmed computer system may comprise any one of a large number of possible software programs that may be designed by those skilled in the art without undue experimentation. Such a programmed computer system comprises a means for determining or designating one or more particular targets from a set of targets to probe (a current target), a means for determining or designating a sequence substring from the current target and determining its complementary sequence (a candidate probe). The means for choosing a sequence substring may function by starting at a particular point in the current target and then incrementing the starting point by some amount each time a new substring is chosen. A substring may be chosen at random or any arbitrary function may be applied in order to choose which substring to pick by the computer means, a means for determining whether the candidate probe satisfies any criteria required or desired for probes, and a means for calculating the T_m for the candidate probe using a hybridization model.

35

40

As used herein, the following terms are understood to mean the following:

A "target" is an oligonucleotide in a sample.

45

A "probe" is an oligonucleotide intended to bind or hybridize to a target. Note that in cases where probes and targets are in solution, a particular oligonucleotide can be both a probe and a target. A "set of probes" is intended to include two or more probes. Preferably, a "set of probes" includes 10

50

55

5 or more probes. More preferably, a "set of probes" includes 100 or more probes. Even more preferably, a "set of probes" includes 1000 or more probes.

10 The "intended target" of a probe is the target that it is designed to best hybridize to. Generally, this will be the target from which the probe is a complementary substring. For example, if GATTACAGATTACA is a particular oligonucleotide in solution (or target), one possible substring is CAGAT. The complement of the substring CAGAT is ATCTG, and thus ATCTG can be a probe for hybridizing to the intended target GATTACAGATTACA.

15 An "unintended target" is a target other than the intended target.

"Cross hybridization" is hybridization of a probe to a target other than its intended target or to another probe.

20 "T_m" is most preferably the melting temperature of hybridization of a probe to its intended target. Melting temperature is defined in scientific literature and is used herein to describe a measure of how strongly a probe hybridizes to a target. More generally, T_m may be any useful measure of the strength of hybridization including, but not limited to, measures such as the best percentage match of the probe against a target, where A matches T and G matches C; the energy of binding of the probe
25 against its target; the negative of the entropy of binding; some combination of the energy of binding and the entropy of binding; the enthalpy of binding; etc.

30 "Cross T_m" is the melting temperature of hybridization of a probe to an unintended target (or to another probe). For melting-temperature models that are location dependent, it is preferable to use the location where the melting temperature of hybridization is highest. Or, as in the description of T_m above, may more generally be some other measure of the strength of hybridization (such as percentage matching, energy of binding, negative entropy of binding, enthalpy of binding, combinations of these, etc.).

35 "Constraints" are the conditions or qualifications that must be substantially met when choosing probes. In most instances, "constraints" refer to a feature or property of the probe. For example, it might be desirable to select only those probes having a T_m between about 50 ° and 60 ° C or that do not contain more than three G's in a row. There may be any number of heuristics or
40 constraints imposed by preferences on the practitioner to use the probe set. Some exemplary constraints include that all probes are a particular length (e.g., 20 bases long or 30 bases long) or that all probes have T_m's within a particular range (e.g., within 5° C of each other or within 2° C of a mean
45 T_m for probes having a particular length). In the case of using probes as primers, there can typically be constraints that the probe must bind to the target sequence within a certain area (in order to do the priming task correctly). Other constraints might be that the probe is not to have G's and C's within the last four base positions of its 3' end, and so on.

5 "delta Tm" is the difference, for a particular probe, between Tm and the maximum cross Tm.
A "hybridization model" is a mathematical model by which one calculates an estimated Tm or
10 cross Tm based on the probe, the oligonucleotide to which it hybridizes and possibly the position of
the hybridization. The hybridization model may also require the input of solution concentrations or
additional factors. An important feature of a "hybridization model" is that it provides an estimate of
Tm or cross Tm algorithmically. There are many hybridization models discussed in scientific
15 literature and are believed applicable within the scope of the methods of the present invention.
Likewise, additional custom hybridization models may be created.

"Acceptable delta Tm" is the smallest delta Tm that is determined to be acceptable for a probe
to be accepted according to the methods of the present invention. For example, an acceptable delta
20 Tm might be 5° C, 10° C or, for greater separation, 20° C. Likewise, an acceptable delta Tm might be
chosen as any number in between. In general, the larger the delta Tm, the easier to separate
unacceptable from acceptable probes. Similarly, the larger the delta Tm, the easier to dehybridize
unintended targets while maintaining the intended targets hybridized.

25 As used herein the term "bind relatively well to intended targets" is understood to describe a
feature whereby a probe does not separate from but rather remains hybridized to a target sequence
under normal operating conditions. A preferred example of such a feature is a perfectly
complementary probe that does not separate from but rather remains hybridized to a target sequence at
30 temperatures under 80° C

As used herein the term "does not bind substantially to unintended targets" is understood to
describe a feature whereby a probe does not hybridize to target sequences other than those to which it
possesses a high degree of complementarity under normal operating conditions. A preferred example
35 of a such a feature is a perfectly complementary probe that does not separate from but rather remains
hybridized to a target sequence at temperatures below 80° C. However, the same probe does not bind
to or easily separates from a target sequence to which it does not possess a high degree of
complementarity at temperatures greater than some temperature significantly below 80° C, such as
40 greater than 70° C, greater than 65° C, greater than 60° C, greater than 50 C, etc. The feature is
intended to include minimal hybridization that may be reversed by agitation or heating above room
temperature.

45 DESCRIPTION OF THE PREFERRED EMBODIMENTS

The following are provided purely by way of example and are not intended to limit the scope
of the present invention.

5

EXAMPLE 1**Selecting oligonucleotides for low cross hybridization**

The following is a sample list of targets for designing probes. The probes were to be built on a DNA chip such as those used in accordance with the method described by Montgomery in WO 98/01221, attached to a layer on the surface of a chip some other substrate so that the probes themselves are not floating around in solution. Thus, we do not account for probes in the target set during operation of the method.

15

>gi|6717738|gb|AW305385.1|AW305385 xv93h12.x1 NCI_CGAP_Brn53 Homo sapiens cDNA clone
IMAGE:2826119 3', mRNA sequence
GCCAGTCACATGCTTACCTGCATTTTAAAGACAGCTTTCAGGTATTTGGGGACTACATTA
TTACCAAAC
CTTGGCTTTGGGAGATTATACAGGTCCGAGGAACTCGTGTCTACTGCAGACGAATGCAAT
TACCCACCT
TCCTCCATACAGAATTGTTAGGAAATGTCCACTCCTTTGGGGTGATTTTCTCCTCAAGT
TGTAGCCAA
CATTTTGTCCGTAAGTATTTCAGGGCAAACATTTCTGACATCTTCTCCAGCTCAGTCTG
CCATGCCTT
GGCAATCCAGTTTCTCCTGTCATATGCGAGCCATCCAAGTTGATGCCAAGTAAGATTTGCCC
AGCTCAAAGT
GAAAGTGTTCGCTCTTGGTATCCGGAATCCTCAGCCCCAGTAGCAAAGCTTTAGTCATTC
ACCTTCATC

20

25

30

>gi|6717590|gb|AW305237.1|AW305237 xr79h11.x1 NCI_CGAP_Lu26 Homo sapiens cDNA clone
IMAGE:2766405 3', mRNA sequence
ACTACTATACGGCTGCGAGAAGACGACAGAAGGGTCATGTGTTAACTATAATCACATTTA
TGGTTTGGA
CCATCACCCCAAGGTAAAAAATAAAGGTATTTCCAGGTATGTTTGGCAAAATAA
AATAAAGGTAA
TTAAAAACCGAAAAAATAAAGGTATTTCCAGGTATGTTTGGCAAAATAA

35

>gi|6714107|gb|AW304418.1|AW304418 xv60f12.x1 NCI_CGAP_Lu28 Homo sapiens cDNA clone
IMAGE:2817551 3', mRNA sequence
CTTTTCTTTTATTCATCTCCAGCAGATCTTCTTTTCTGTAAGCTTACCACTTCTAAA
TTTAATAT
GTGTTTTGAGCTCATTATTTAAAGGAATCACATCTTGCTAATCACATCCAAGGCACCGGA
ACATAGTGTC
TATACTGACTGAACAGGCCAAGCTTCGTGAGTTAATTAATAAAATATTTGGTAAGAAACG
GTCCATCATT
ATCTTATCACTTGAGATGACAATGTTGAAACTTACAGGATGGAAGGCATCTCATTAAATC
AGACCATTTT
AAATCAATTTTATTTGACTTACAGTCTTGAAATAACATATCATTATCTTTGGCCATATCA
AAAAGTGA
CCCAGTTGGAAAATATTTATATGTCCAAATATTGGTTTAGAGGAAAGTATAGCATGTTTTT
GGTAAAT

40

45

50

55

5

>gi|6713707|gb|AW304018.1|AW304018 xv15h11.x1 Soares_NFL_T_GBC_S1 Homo sapiens cDNA clone IMAGE:2813253 3', mRNA sequence

10

TTTTTACAGGATAATACITTAATTACGAAAAGCACAAATTATGTATCATTGGAACCAGC
AAACACACAG
TAGCAGGAAGGATGCTTGCTCCCAAGGCTCTTCAGTCATCAGAGGACACACTCAAGCCCC
ACCTGAGTCT
TCTCCCCATTCCATCGGCCATCCCTGCTCAGGATGTGGTACCAGGGCCATTCCCAACAGCC
TCATCTCAG
TAGACTCCAGTTTGTCTAATTCCTCTCAATGGTGCTCCTTCGTCACCTCTCGTGGGCTGGC
GGATAG

15

>gi|6713507|gb|AW303818.1|AW303818 xr23d05.x1 NCI_CGAP_Ut4 Homo sapiens cDNA clone IMAGE:2760969 3', mRNA sequence

20

GAACAGGAA
TAATAATTTACAAATACTAACACTTTATTGACAATAGACAAGTCTTTAGGGTAGTGCA
CATGTACTTA
AAAACTACCTTCTACCAATCTCAACACTTTTATAAATTTTCAGGTGAAACTGTAGCAGAT
CCTACTTTA
TTTTTCAATGGTTAGTGTAATAATTCTGTATGTAAAATAAGTACATATTTTGAGATGGAAGA
AGGACTGCA
TGTGAAATGCTTTGCCTAAGTTGTAAGGCTCCTGTCTTTACGCTATCATTAAAGGCCAAAA
AAATCACTG
CTAGAAATGTTCCCCAAAAAATCTTAAACAGCTCAGTCTTTAAAAGTATTAATAATTTTT
TTTTTTTTT
TTTTTTGGAGACAGAGTTTCGCTCTTCTTGCCCAGGCTGGAGTACAATGGCGCAATCTCAG
CTCACCG

30

>gi|6712898|gb|AW303218.1|AW303218 xr59g03.x1 NCI_CGAP_Ov26 Homo sapiens cDNA clone IMAGE:2764468 3' similar to contains Alu repetitive element; mRNA sequence

35

TATACGGCTGCGATAAGACGACAGAAGGGGTAGGACTGAGGCCTGAGTACACCTTTTAT
ATTTTGGACAT
TTACGTATTAATAAAAAATTATCTAGCTGGGCATGGTGGCACACACCTATGGTCCCAGCTGC
TTGGGAGGCT
GAAGTAGGAGGCTGGCTTGAGCCCAGGAGTTTAAGTCCAGCCAGAGCAACATAGTGAGA
ATTCATCTCAA
GAAAAAAAAAAGAAAAAAAAAAAAAAAAAGAAAAAAAAAGTCGTATCGA

40

>gi|6711895|gb|AW302218.1|AW302218 xs03d05.x1 NCI_CGAP_Kid11 Homo sapiens cDNA clone IMAGE:2768553 3' similar to TR:Q14934 Q14934 NF-AT3. ; mRNA sequence

45

CATATTACTGGTCATTGAGCAGTTTATTGGGAGCAATCTGACCCAGGTTGCCAGCACAA
CAGCCAGCCC
ACACTCTAGACACGCCTTCACTCCAGTCCATTCTGGCACCTAGCCTCAGTCTTCAACCTCC
TCCCTCCTC
CACACACTCCTTCCCCAGCCCTCCAAGGCAGCACCAGGCCTGAGGGCCACACCTCAGCT
GGGGGAGGGG
AGGGAAGACAGTGAGACAGACAGAAGCTGGGGAAAGAGGAGCCAGGGTTGGCCCCAAG
CTTCTGTAGCCA
CCACTCCAGGAAGGAGGGAAAGGGGGCAGGGCTGAGGCTGGGGCTGGGGTTGCCAGGTG
ATGACAGTTCA

50

55

5

CGTGGTTCAGGCAGGAGGCTCTTCTCCAGGAGGTGCAGGGAAGCCACTCAGGTCTCGGCC
AATGATCTCA
CTCACTGTAAAGGAGGGGCAGTTGAGAGACTGGGCTA

10

>gi|6710695|gb|AW301018.1|AW301018 xk11e01.x1 NCI_CGAP_Co20 Homo sapiens cDNA clone
IMAGE:2666424.3', mRNA sequence

GCCCTTTTTTTTTTTTCAACAAGGAAATGTATTTATTTTTCTTTAGAATTTGGCTCAG
AACAGTAA
CAAAAATATTTACATTAAAATAAAATTAACATGCAATTACTTAACCATATGTAATAATTTA
CGTTGGAATA
TATTAGCCTTCCCATGAGTTTAATAAAAACTAATATTTGGTTTTAGATTCAATACCATCCT
TTCAAATAT
TTGGGTATGAACTTGGTAGCAATGCAATTGTCTGATGTACAGAGCAGATTTACCATGA
GAGATTACAC
CAAAGAACAGATGTCCCTTCCAGAACATTATCTACCCCAGACTCAGAACTGAGCAGC
CAAGCTTCCT
TCCCAGGAATCACCATGGAATGTCTGAACAATAACCAGGCCCTGGAGATTACTGCAGGGC
TGCCAGAGTT
TTAGGAATCAGCCAAACTC

15

20

25

>gi|6710495|gb|AW300818.1|AW300818 xk06e09.x1 NCI_CGAP_Co19 Homo sapiens cDNA clone
IMAGE:2665960.3' similar to TR:O88814 O88814 HEME-BINDING PROTEIN. ; mRNA sequence

AAGTCAATGCCTTTATTTTTAGTTTTCTGAAGACAAAGCTCTTATAAGAATCACAGATG
AAAGATCAG
GCACAAATCACATTTTCCCCCTTAATAACAAAATACAAATCCAATAATTTAGAAAATCA
GTTTTTAGTG
ACCCAGATGCCTGGAGAAAAAGCTGCCAGGATTTTTCTGGTCTATCGCAGAATTTCTACA
TCAATGAGAA
GGATGCTGCATATCTTGGCTGTATTATTTCTACCGTGAGAAAAGAGACTTAGTATATGG
AACATGCTTT
TTTCAGAAAATTGGCAGTAAGTACCTTTGAAGGAAAGTTGGTTAAGTTGGAAGTTGCAGCT
GGAAGTTGGG
AAGCACTGTCCCTCCTTACCCCGAGGAAGGAGACACAGAGGCACACTTCCAGTAAGTT
CTTGGTTCAG
TGGGTCACTCATGTCTTCAACAGCCAGATCTCATTGCGCCGTCCTAGGGCTTCATGGGA
GGGTACATAAC
CCGTGCAGAAATAGATGTCCNCCGGTAGGGTGCTGTGCCCTTCAAGGGAGCACGCAA

30

35

40

>gi|6705458|gb|AW298822.1|AW298822 UI-H-BW0-ajq-h-09-0-UI.s1 NCI_CGAP_Sub6 Homo
sapiens cDNA clone IMAGE:2732800.3', mRNA sequence

CGGCCGCGCCGGTTTTTTTTCAAGTTTTGGGTATGTTAATCTGTTATGTACTACTGTCTGT
TTTGTAT
TGTTTTGTAAATTACACCATAATGCTAATTTAAAGAGACTCCAAATCTCAATGAAGCCAG
CTCACAGTGC
TGTGTGCCCCGGTCACCTAGCAAGCTGCCGAACCAAAAGAATTTGCACCCCGCTGCGGGC
CCTTGGTT
GGGGCCCTGCCCTGGCAGGGTCATCCTGTGCTCGGAGGCCATCTCGGGCATAGGTCCACC
CCGCCCCACC
CCTCCAGAACCGGCTCACGCTTACCTCAACCATCCTGTTTGCGGCGTCTGTCTGAACCAC
GCGGGGGCC

45

50

55

5
 TTGAGGGACGCTTTGTCTGTCGTGATGGGGCAAGGGCACAAGTCCTGAATGTTGTGTGTA
 TCGAGAGGCC
 AAAGGCTGGTGGCAA

10
 We used the following constraints. Probes must be 20 bases long. Probes must have a T_m within 1°C of the expected T_m for a 20-mer according to the hybridization model used (in this case 68.25°C).

15
 We used the following hybridization model: $T_m = 81.5 + 0.41 * P_{gc} - 675 / N - P_{mm}$, where P_{gc} is the percent GC content of the probe = (number of G's + number of C's) / $N * 100$, N is the length of the probe in bases, and P_{mm} is the percent mismatches = (number of mismatches) / $N * 100$.

We chose an acceptable delta T_m of 20°C .

20
 The algorithm worked as follows. We began with target 1. We picked a 20-mer out of it at a randomly selected location and found its complement as a candidate probe. We checked that the candidate probe satisfied the constraints. If not, we chose another 20-mer from a random location. If it did, we then calculated the T_m 's for this probe hybridizing to all other targets at all other locations and used that data to find the maximum cross T_m and thus delta T_m . If delta T_m was greater than or
 25
 equal to 20°C , we kept this probe and obtained a probe for the next target (target 2). If not, we chose another 20-mer from a random location. We repeated this process until we found one acceptable probe for each target.

30
 The following is the list of probes found by this process. In the following, there is header information given for each probe indicating from which target it comes (i.e., what its intended target is), where in that target the probe comes from (i.e., at what offset into the intended target), the T_m of the probe, the maximum cross T_m of the probe, what unintended target provides the maximum cross
 35
 T_m , and where in that unintended target the maximum cross T_m happens (at what offset).

Note that the T_m values given all match exactly. The experimentally determined T_m 's will not necessarily match exactly -- the T_m 's given are estimated T_m 's derived from the hybridization model, which in this case results in the methods described being able to find probes that all match
 40
 exactly in estimated T_m .

45
 > probe 1 from target 1 at offset 359; $T_m = 68.3$; max. cross $T_m = 33.3$ from target 9 at offset 253
 ATTCCGGATACCAAGACGCA
 > probe 2 from target 2 at offset 2; $T_m = 68.3$; max. cross $T_m = 48.3$ from target 6 at offset -3
 CTTCTCGCAGCCGTATAGTA
 > probe 3 from target 3 at offset 145; $T_m = 68.3$; max. cross $T_m = 28.3$ from target 5 at offset 272
 AAGCTTGGCCTGTTCACTCA
 > probe 4 from target 4 at offset 239; $T_m = 68.3$; max. cross $T_m = 23.3$ from target 1 at offset 176
 AAGTGACGAAGGAGACCAT
 > probe 5 from target 5 at offset 424; $T_m = 68.3$; max. cross $T_m = 28.3$ from target 1 at offset 25
 50

5

GAGCGAAACTCTGTCTCCAA

> probe 6 from target 6 at offset 36; Tm = 68.3; max. cross Tm = 28.3 from target 1 at offset 225

AAAGGTGTACTCAGGCCTCA

> probe 7 from target 7 at offset 333; Tm = 68.3; max. cross Tm = 28.3 from target 1 at offset 314

10

ACGTGAACTGTCATCACCTG

> probe 8 from target 8 at offset 352; Tm = 68.3; max. cross Tm = 28.3 from target 5 at offset 24

CATTCCATGGTGATTCTCTGG

> probe 9 from target 9 at offset 210; Tm = 68.3; max. cross Tm = 33.3 from target 4 at offset 79

AGCCAAGATATGCAGCATCC

> probe 10 from target 10 at offset 350; Tm = 68.3; max. cross Tm = 28.3 from target 1 at offset 175

15

ACAGACAAAGCGTCCCTCAA

EXAMPLE 2

20

In this example, the list of targets was the same as in Example 1. Likewise, all of the parameters and the model used for calculating Tm were the same as in Example 1. The only difference was that we used a variation of the method.

25

We began with target 1. We picked a 20-mer out of it at a randomly selected location and found its complement as a candidate probe. We determined that the candidate probe satisfied the constraints. If not, we picked another 20-mer from the "next" location, *supra*. If it did, we calculated the Tm's for this probe hybridizing to all other targets at all other locations and used that data to find the maximum cross Tm and thus delta Tm. If delta Tm was greater than or equal to 20° C, we kept this probe and moved on to getting a probe for the next target (target 2). If delta Tm was not greater than or equal to 20° C, we went back and picked another 20-mer from the "next" location (see below). We repeated this process until we found one acceptable probe for each target.

30

35

By "next location," we applied the following process. We selected a new candidate probe starting at a location one base to the right (in the 3' direction) of the previous pick. If such a location resulted in not having enough bases to make a candidate probe (such as when the next location is too close to the end of the target so that there are not enough bases left to make a probe of the desired length), we started at location 1 of the target. Thus, the process of scanning a target for an acceptable probe was started at a randomly selected point and then progressed incrementally along the target with wrap-around to the front of the target when the end was reached.

40

45

This process provides an exhaustive search of a target for an acceptable probe. It will find an acceptable probe if one exists. Thus, it is a good candidate search method for situations where the targets might be very similar except for small differences (perhaps mutations) at particular sites in the oligonucleotide.

This process resulted in the following set of probes being found.

50

> probe 1 from target 1 at offset 62; Tm = 68.3; max. cross Tm = 33.3 from target 3 at offset 372

55

5

CCCAAAGCCAAGGTTTGGTA

> probe 2 from target 2 at offset 0; Tm = 68.3; max. cross Tm = 38.3 from target 6 at offset -5

TCTCGCAGCCGTATAGTAGT

> probe 3 from target 3 at offset 115; Tm = 68.3; max. cross Tm = 33.3 from target 6 at offset 173

10

TATGTTCCGGTGCCTTGGAT

> probe 4 from target 4 at offset 234; Tm = 68.3; max. cross Tm = 33.3 from target 1 at offset 399

ACGAAGGAGCACCATTGAAG

> probe 5 from target 5 at offset 292; Tm = 68.3; max. cross Tm = 33.3 from target 7 at offset 111

GGAGCCTTACAACCTTAGGCA

> probe 6 from target 6 at offset 154; Tm = 68.3; max. cross Tm = 33.3 from target 3 at offset 73

15

TTAAACTCCTGGGCTCAAGC

> probe 7 from target 7 at offset 265; Tm = 68.3; max. cross Tm = 28.3 from target 4 at offset 93

AGTGGTGGCTACAGAAGCTT

> probe 8 from target 8 at offset 379; Tm = 68.3; max. cross Tm = 28.3 from target 1 at offset 59

ATCTCCAGGGCCTGGTTATT

> probe 9 from target 9 at offset 438; Tm = 68.3; max. cross Tm = 38.3 from target 4 at offset 194

20

CGCAATGAGATCTGGCTGTT

> probe 10 from target 10 at offset 350; Tm = 68.3; max. cross Tm = 28.3 from target 1 at offset 175

ACAGACAAAGCGTCCCTCAA

25

Although the invention has been described with reference to the presently preferred embodiments, it should be understood that various modifications can be made without departing from the spirit of the invention. Accordingly, the invention is limited only by the following claims.

30

35

40

45

50

55

Claims

5

10

15

20

25

30

35

40

45

50

55

5

WHAT IS CLAIMED IS:

10

1. A method to provide a set of probes that hybridize relatively well to their intended targets but do not substantially hybridize to unintended targets comprising the steps of:

15

- (a) Determining a set of targets;
- (b) Determining a particular current target from the set of targets to probe ;
- (c) Choosing a sequence substring from the current target and providing its complementary sequence, which becomes the candidate probe;
- (d) Determining that a candidate probe satisfies any criteria desired or required for probes;
- (e) Calculating the T_m for the candidate probe using a hybridization model;
- (f) Calculating substantially all possible cross T_m 's of the candidate probe hybridizing to all unintended targets and finding the maximum cross T_m ;
- (g) Calculating delta T_m ;
- (h) Determining whether the delta T_m is acceptably large.
- (i) Repeating steps (b) forward until the desired probes are found.

20

25

30

2. The method of claim 1 wherein choosing a sequence substring is performed by starting at a particular point in the current target and then incrementing the starting point each time a new substring is chosen by some amount.

35

3. The method of claim 1 wherein the substring is chosen at random.

4. The method of claim 1 wherein the delta T_m is at least about 20° C.

40

5. The method of claim 1 wherein the delta T_m is at least about 10° C.

6. The method of claim 1 wherein the delta T_m is at least about 5° C.

45

7. A set of probes that hybridize or bind relatively well to their intended targets but do not bind substantially to unintended targets.

50

8. The set of probes of claim 7 wherein the delta T_m of the set is at least about 20° C.

55

5

9. The set of probes of claim 7 wherein the delta Tm of the set is at least about 10° C.

10

10. The set of probes of claim 7 wherein the delta Tm of the set is at least about 5° C.

15

11. A set of probes that hybridizes or binds relatively well to intended targets but do not bind or substantially hybridize to unintended targets produced in accordance with the method of claim 1.

20

12. A programmed computer system for providing the sequences of a set of probes that hybridizes or binds relatively well to intended targets but that does not substantially hybridize or bind to unintended target comprising a software program having a means for determining or designating one or more particular targets from a set of targets to probe (a current target), a means for determining or designating a sequence substring from the current target and determining its complementary sequence (a candidate probe), a means for determining whether the candidate probe satisfies any criteria required or desired for probes, and a means for calculating the Tm for the candidate probe using a hybridization model.

25

30

35

40

45

50

55